



Cover Crops

A practical guide to soil
and system improvement



In association with

Kellogg's
ORIGINS



Kellogg's Origins™
Natural Heritage

Introduction

This NIAB TAG and Kellogg's Origins™ publication is a practical guide to the use of cover crops for soil and system improvement. It provides information on the selection and management of cover crop options. Cover crops can (among other things) help to increase resource use efficiency, improve system management/resilience, benefit environmental goals and increase yield/economic returns. Origins™ farmers are working with NIAB TAG to translate research into practice and to quantify the delivery of key benefits.

The guide provides information based on current understanding on how to select, deploy and make the most of specific cover crops against identified end uses. Guidance has been generated from a range of sources including research, grower feedback and other expert opinion. There may be a need to amend specific practices for individual farm situations.

It also serves to support and inform on-farm decision making and will be developed further in collaboration with field activities and feedback from the Origins™ farmers.

The focus is primarily on the use of autumn sown cover crops used ahead of spring sown crops, but also features some aspects of cover crop use in wider scenarios. To use the document select the cover crop goal which best describes your objective; then use the decision guides to select an appropriate cover crop species or mix. The cover crop options are suggestions and further ingredients or particular agronomy and management inputs may be required to help tailor the selection to a particular circumstance on farm.

Ron Stobart

Ron Stobart

Head of Farming Systems Research, NIAB TAG
E: ron.stobart@niab.com



This publication may not be reproduced in whole or in part, stored in a retrieval system, transmitted or circulated by electronic, mechanical, photographic or other means without the prior permission of NIAB.

While every care has been taken in the preparation of the advice contained in this booklet, NIAB TAG and Kellogg's cannot accept responsibility for any loss of inconvenience arising from following the information herein.

This edition published August 2015
© NIAB Cambridge 2015
A charitable company limited by guarantee

Designed and produced by Cambridge Marketing Limited, 01638 724100

Contents

Introduction	inside front
Costs and returns	2
Case study	4
Selecting your ingredients – choosing your cover crop	5
Managing weeds and pests	5
Environmental goals	7
Soil fertility building and nutrition	9
Improving soil structure	9
The ingredients – cover crop components	14
Brassica types	14
Legume species	16
Other options	19
The mixing bowl – combinations of components in a cover crop	21
‘Cooking’ – growing the cover crop	25
Seed sourcing	29
Further information	29

Costs and returns

Cover crops can provide benefit across the rotation (e.g. mitigating pollution, reducing erosion and improving soils/habitat), but direct financial benefit is most easily gauged against yield improvement in following crops. While some variation is expected, the following is a provisional guide based on research, farm feedback and expert opinion.

This section gives an indication of both yield and potential margin changes that would be needed to deliver a financial benefit.

Response

Benefits from cover crops accrue through improvements in nutrient cycling, water retention, soil structure, soil biology and other parameters. Finding positive changes in these areas is a useful indication of a direction of travel and is often detectable before yield responses. However, yield is the best absolute measure and is often seen in crops following cover crops, but sometimes cannot be detected until later in the rotation (or responses may require repeated cycles of cover crop use to become fully apparent).

Yields

Findings from longer term studies at NIAB suggest:

- Yield response in the crop following the cover crop will vary with cover crop, following crop type, season and management. Some following crops have shown little response and others more consistent yield improvements.
- Research examining the yield response in spring barley following an autumn legume cover crop was assessed in NIAB TAG National Agronomy Centre research at Morley in Norfolk over four years (i.e. the trial was repeated in each of four seasons). While responses in individual seasons varied, a mean yield responses of c. 0.3 t/ha was seen in spring barley following use of a legume cover crop. This response was achieved with a 'farm standard' nitrogen (N) dose (of 150 kg/ha) applied to the spring barley.
- Longer term NIAB TAG New Farming Systems research at Morley (running since autumn 2007) has also shown responses in wheat crops behind the crop following the cover crop (i.e. a rotational scenario of autumn sown cover crop, spring sown break crop and then autumn sown winter wheat). The response will again vary with cover crop type, season, repeated use and management; however, studies have suggested yield responses of the order of 0.25 t/ha (at standard N doses) following legume and brassica cover crops can be detected.
- Linking these research findings together gives a potential cumulative grain return of c. 0.5 t/ha over cereal crops following the use of a cover crop.
- In addition to the direct financial benefits, cover crops can also provide wider value on farm (e.g. reduced soil erosion risk and improved surface drainage) and potentially contribute to longer term 'climate smart' improvements to soil (e.g. improved workability, moisture retention and soil organic matter) and the overall farm system (e.g. greater resilience).



Grain value

The value of 0.5 t/ha of grain depends on the mean grain price over the two crops. Examples of value against a range of prices are set out in Table 1; this indicates what cover crop cost would cover the gain

(i.e. a break even point). Table 2 sets out the yield increase needed to break even at a range of potential cover crop costs.

Table 1. The value of grain yield margin response (£/ha) for a range of mean grain prices (£/t) at a series of anticipated yield benefits (t/ha).

Grain value per tonne (£)	80	100	120	140	160	180
Yield benefit expected (t/ha)						
0.3	24	30	36	42	48	54
0.4	32	40	48	56	64	72
0.5	40	50	60	70	80	90
0.6	48	60	72	84	96	108
0.7	56	70	84	98	112	126

Table 2. Yield response required (t/ha) for a range of sample cover crop costs (£/ha) at specimen grain prices (£/t).

Grain price (£/t)	80	100	120	140	160	180
Sample cover crop cost (£/ha)						
20	0.25	0.20	0.17	0.14	0.13	0.11
40	0.50	0.40	0.33	0.29	0.25	0.22
60	0.75	0.60	0.50	0.43	0.38	0.33
80	1.00	0.80	0.67	0.57	0.50	0.44
100	1.25	1.00	0.83	0.71	0.63	0.56

Costs

These can vary substantially but, in work undertaken at The GWCT Allerton Project in Leicestershire and at NIAB, seed costs have varied from £15-60/ha. Costs of establishment and management systems also vary and should be guided by condition and requirement. Establishment systems range from £15-30/ha for broadcasting/direct drilling based systems to over £40/ha for combi-drilled systems.

Input costs are generally low but may range from nothing to £30/ha including application costs (e.g. where starter fertiliser and pest protection is needed). Growers should calculate sample costs based on their own systems, but specimen costs would often be £50-70/ha with a potential range of £30-130 at the system extremes.

Wider benefits

The financial returns listed here do not include any wider benefits (soils, weeds, habitat, soil organic matter (SOM) etc). While there is a clear benefit, for example, to improvements in soil organic matter, the financial implications of this are difficult to resolve and are likely to vary with season and circumstance. In addition, financial benefits against the use of cover crops can also be accrued through environmental schemes; the additional benefits are also not included in these calculations.

Case study

Kellogg's Origins™

Phil Jarvis is Head of Farming for the Game and Wildlife Conservation Trust's Allerton Project, based at the Loddington Estate in Leicestershire. The estate is also a LEAF Innovation Farm and the Defra SIP mixed arable and livestock Study Farm.



Size	319 ha – as part of a 900 ha joint farming venture
Soil	Clay soil (Hanslope and Denchworth series with a small amount of Banbury)
Cropping	Winter beans, winter wheat, winter oilseed rape, winter and spring oats
Livestock	280 mule ewes and, through a grazing agreement with a neighbour, a South Devon suckler herd
Staff	Two staff
Machinery	NH 9080 Combine, Caterpillar 765, 7530 John Deere tractor, Dale 4m eco drill, Kuhn fertiliser spreader

The Allerton Project, in association with the Game and Wildlife Conservation Trust (GWCT), was set up in 1992 to carry out public, private and charitably-funded research on the interaction between farming, the environment and wildlife, and to advance industry and public education on these issues. A commercial farming operation runs alongside research and demonstration studies on the effect of farming systems on wildlife, soil conservation, water and environmental habitats.

Since 1992 the soil management strategy on the farm has moved progressively toward non inversion tillage and more recently single pass (direct) drilling. The next chapter in the farmland research includes resource protection, reducing fuel and energy consumption and increasing soil resilience and organic matter with the aim of working towards a sustainable farming blueprint. This means maximising the value of agrochemical inputs and using more biological inputs from natural processes; our move to include cover crops in our rotation encompasses many of these points. We currently grow oil and tillage radish with some oats, but have been experimenting with other mixes as well.

The cover crop captures nutrients, making them available for following crops and the soil; this improves the soil structure, reduces our erosion risk and there is some evidence that the right sort of cover crop may help with our overall black-grass management strategy. We are following this up with a range of practical trials and demonstrations on farm particularly through the Kellogg's Origins™ project, but also other initiatives such as the Defra SIP platforms.

However, from our experience growing cover crops is not a quick and easy option. When planning a cover crop, especially if direct drilling, we have learnt to take into account the previous cropping and spray regime, particularly with sulfonylureas such as Atlantis which can have a residual effect. Slugs, trash from the previous cropping and dry conditions can also reduce establishment. You also need to give thought to cover crop destruction and the establishment systems for your following crops.

The economics must also be considered. We often talk about soil resilience, with a robust rotation and soil biology, but we're in the business of profitable food production. Interest in cover crops will be dictated by the gross margins, fixed costs and general profitability of the system. Our aim is also to put some costs on these farming systems; are the additional establishment costs, including an extra drilling, affordable and worthwhile and how does the cover crop compare to other methods of intervention, for example agrochemicals for weed control? Our trials include different crop species, varieties, and seed rates with costs ranging from £20 to £50/ha. We want to look at the impacts of these approaches in the following crops and across the rotation so we can get a better idea of the value as well as the cost.

Selecting your ingredients – choosing your cover crop

Managing weeds and pests

The use of cover-crops to help manage weed and pest populations is a developing area; it is in need of further research and may not be suited to all cover crop circumstances. This section will focus ostensibly on weed management scenarios, but will include detail on pest management where appropriate.

There are a number of potential approaches to the management of grass-weeds through the use of cover crops, but essentially, cover crops can be used for short term benefit or longer term strategic weed management; these systems are not necessarily mutually compatible. AHDB funded research (RD-2012-3789) being delivered through Rothamsted and NIAB examines the relative merits and interactions of these options in greater detail.

For longer term strategic management the cover crop is fundamentally a trap-crop; the pest, in this case black-grass, is encouraged to establish and thrive, and then, before the pest can reproduce, the pest and the cover crop are destroyed together. The grass weed population is reduced over time by a combination of seed removal from the seed-bank, germination and establishment in the cover crop, and natural process of seed death in the soil.

For a shorter term approach, the cover crop can essentially be used to suppress weeds ahead of a following crop. While this can deliver benefit to a following spring crop (perhaps where herbicide

options may be more limited) any effect of the cover-crop on reducing the number of black-grass plants that germinate and establish, potentially leaves a higher number of seeds in the seed-bank, to carry over to the following crops in the rotation.

Current research is comparing the effectiveness of the underlying cropping strategy (e.g. the inclusion of spring cropping or fallows) with the effectiveness of the same system including various cover-crop approaches. This will enable the effects of the cover cropping approaches, over and above the underlying strategy, to be demonstrated and quantified. While research remains ongoing early indications and field experience perhaps suggest that the underlying system changes, along with the wider merits of cover crop use, are delivering the more consistent benefits.

With regard to cultural control of black-grass; if the strategy is to exhaust the seed bank prior to planting a crop (whether that is a spring crop or an autumn drilled crop following a year-long fallow) do not use deep cultivation that will mix the soil profile and reduce the effectiveness of that strategy. If spring cropping is the underlying approach and grass population levels have become a real challenge, pick a spring crop where the weed can be managed effectively. Use of non-selective weed control, both to destroy the weeds before they can set seed and (if relevant) to remove any weed seedlings that are present before the crop is drilled is essential.

A cover crop can act as a trap crop for grassweeds such as black-grass, but both pest and cover crop must be destroyed before seed dispersal

Managing weed populations (weeds)		Sanitising cover crops (weeds and pests)		Other biological routes (weeds and pests)	
Trap crops (weeds)	Crop competition	Bio-fumigation	Trap crops (pests)	Allelopathic effects	Habitat creation (pests)
A cover crop that facilitates weed establishment and is then destroyed before the weed can reproduce.	Cover crops that outcompete weeds can help to provide a clean seedbed for the following crop.	Some brassica species have high levels of isothiocyanate; this can sterilise soil. Such cover crops can be used against weeds and soil pests.	Some cover crops (notably brassicas) can promote egg hatch in some pest species e.g. this can be effective against certain nematode types.	Some cover crops (e.g. clovers, rye and oats) can have allelopathic activity; inhibiting the germination of weed and other species.	Pest management can also be delivered through improving predator habitat.
Suggested options/ingredients					
A wide range of cover crops can be used for this purpose. Mixtures are common and components could include brassicas cereals or legumes.	Brassica species (e.g. radish and mustards) that can cover the ground are common; but a range of cover crops could be used.	Choose a specific variety (usually radish or mustard) that is sold for this activity. Research is ongoing in this area and there is a need for more impartial comparative data between types.	Select a variety sold for this activity. Variety choice should be guided by recent research based comparisons (e.g. current levy body research outputs as available).	Limited current UK field information and further research is needed in this area. Consult specialist cover suppliers for options.	Potential to use cover crops strategically on farm to provide habitat or companion crops for wide row species.
Other comments (also see agronomy guidance pages for specific suggestions)					
This approach is used frequently and mixes need to be open enough to allow weed germination. Origins™ research has also shown starter fertiliser can increase weed populations in cover crops.	Rotation conflicts can be an issue where brassicas are used. Give particular attention to destruction (to remove weeds in the cover crop) and minimising soil disturbance when drilling to maintain benefits accrued.	Methods of production, destruction and incorporation are important; follow seed supplier guidelines.	Methods of production, destruction and incorporation are important; follow supplier guidelines.	Academic research has shown effects but there is little impartial field information on magnitude or comparison between types. Effects can be variable and potentially may impact on weeds and following crops.	Data and farm experience supports the value of predators and French companion crop data has shown changes in pressure with some pests in OSR. Further UK based research is needed in this area.

Environmental goals

Habitat creation: Cover crops provide winter cover and a habitat for birds, mammals and insects as well as grazing opportunities for livestock and wildlife. Predator habitat generation through cover crops can also potentially augment pest management.

A wide range of cover crops can be used for wildlife habitat provision, although some may be better suited to specific goals than others. Selection can be guided through expert advice (e.g. via GWCT www.gwct.org.uk).



Cover crops provide winter cover and a habitat for birds, mammals and insects

Nutrient loss: A range of research projects have shown that nitrogen (N) uptake by cover crops reduces the risk of nitrate leaching losses over winter. Reduced soil run-off also lessens the loss of phosphate attached to soil particles. Recent research carried out on Origins™ sites suggested N leaching reductions of 25-54% (mean c. 40% or c. 38 kg/ha N); this N will therefore be retained to benefit crops and the wider soil system (Table 3). A range of cover crops are suited to this purpose, but fast growing non-legume species (e.g. brassicas) are used typically.

Table 3. Nitrogen (N) retention at six Origins™ cover cropping sites in spring 2015. Comparing mean cover crop values to a control area without a cover crop; N values are based on soil N (0-60 cm assessed spring 2015), N in the cover crop foliage (assessed spring 2015) and N in any weed biomass (estimated from GAI).

kg/ha N	Site 1	Site 2	Site 3	Site 4	Site 5	Site 6	Mean
Nitrogen – cover crop system	150	96	71	74	74	77	90
Nitrogen – no cover crop	69	39	44	44	54	59	52

Soil erosion and water quality: Cover crops provide ground cover during risk periods for soil erosion by wind and/or water. Losses are more likely to occur in autumn and winter when soil moisture and rainfall levels are high. Reducing these losses can benefit the environment by lessening losses of sediment, soil, nutrients and pesticides, which can impact on water quality and habitats. Cover crops help to reduce erosion losses mainly through the provision of ground cover at critical risk periods.



Cover crops can help to reduce erosion losses

As a general guide, once more than a third to one half of the ground is covered, there is a substantial reduction in run-off and erosion risks. Species mixtures, such as oats and brassicas, which grow rapidly in the autumn and provide good ground cover, can be a useful simple tool in such scenarios to help reduce erosion risk. Other mixes can be used to suit circumstance, as long as they provide the requisite ground cover. These mixes can also often be used to achieve other objectives (e.g. habitat).

Other considerations: Think about potential conflicts with other crops in the rotation and the management of volunteers in following crops. Where wind erosion is the main issue a cereal cover crop or companion crop (sown with the main crop and then destroyed) can be beneficial. Also think about other methods of erosion management; for example, tramline management (to reduce direct water flow) or strip tillage (leaving stubble between the rows) can also reduce erosion risk.



Chopping a mustard cover crop

Soil fertility building and nutrition

Cover crops can be used as green manures to add organic material back to the soil. This will help to stimulate and feed biological activity in the soil and, in the longer term, regular use of cover crops can raise soil organic matter content. In the NIAB TAG

New Farming Systems programme the use of fertility building cover crops has demonstrated rotational yield and margin (over nitrogen, N) improvements from the use of specific approaches.

Table 4. The impact of cover crop on worm numbers compared to a farm standard or stubble areas over four Origins™ sites (spring 2015). Data presented as a percentage of the number of worms in the cover crop area. Worms sampled from a surface level 20 cm³ block and numbers were: site 1 (115 per m³), site 2 (338 per m³), site 3 (663 per m³) and site 4 (375 per m³).

	Site 1	Site 2	Site 3	Site 4	Mean
Cover crop (%)	100	100	100	100	100
Farm standard autumn cultivation	43	111	–	70	75
Stubble area	–	59	66	–	63

Field evaluation carried out on Origins™ farms has also shown the use of cover crop to improve earthworm numbers; a useful indicator of soil biological activity (Table 4).

Cover crops can also help capture N; this is then potentially available to both following crops and other elements of the soil system and biology. The amount of N released by a cover crop depends on a range of factors, but is influenced by cover crop type, growth, the environment as well as C:N ratio (the carbon to nitrogen ratio in the material) and other biological compounds. Broadly, if the C:N ratio is less than c.13, net mineralisation is observed and N can become available to plants, but as it increases this availability reduces; as the ratio extends past 26 a net immobilisation can occur (for example N is retained elsewhere in the systems by micro-organisms). The extent to which N contributes to crop yield depends on time of the release as well as amount; this can be too early (may contribute to growth but not yield) or too late (could increase grain protein or still give a yield response in later crops in the rotation).




Cover such as fodder radish can help improve the soil structure

Improving soil structure

Cover crops with a vigorous and active root system can help open up soil, to improve structure. Field evaluation carried on a range of Origins™ farms has shown improvements in both soil structure and reduced variability following the use of cover crops.

Findings from one site (Figure 1) indicated, in a stubble area, structure remained similar over winter, but that cover cropping improved structure to a similar level to that achieved with an autumn cultivation (potentially lessening the need for the cultivations). When selecting a cover crop for improving soil structure it is important to assess the structural issues and know the extent and depth of any impediments.

Figure 1. The impact of a range of cover crops compared to an autumn 'farm standard' cultivation on VESS (Visual Evaluation of Soil Structure) pre and post winter on a Kellogg's Origins™ site 2014/15 (a lower VESS score is indicative of a better soil structure).



The following key highlights potentially suitable options.

SOIL FERTILITY

KEY TO SELECT A SUITABLE COVER CROP/MIX COMPONENT

Green manure	Nitrogen (N)	Phosphate (P)	Other nutrients	Improved rooting
Range of cover crop types are suitable as 'green manures' to trap N and add organic material to the soil.	A range of legume species can be used to fix N and in many cases also improve soil structure.	Potential from some polygonums (P scavengers) and legumes (with P cluster rooting).	Cover crops with active rooting can potentially help mine and cycle nutrients.	Improved ability of roots to explore the soil can be of benefit (e.g. lower critical P values).
Suggested options/ingredients				
Oats, phacelia and brassicas (such as mustard or radish) and legumes (faster growing species sown early are more likely to fix N).	Black medick, a range of clovers (e.g. crimson clover), as well as other legumes such as vetch and lucerne; use of species mixtures often well suited here.	Buckwheat and lupins are possible options. Certain mycorrhiza can also benefit some species (not brassicas).	Consider species with extensive root systems or mixtures with complementary rooting.	See section on improving soil structure.
Other comments (also see agronomy guidance pages for specific suggestions)				
Growth and biomass are important, but consider incorporation issues. Fast growing species also tend to trap more autumn N.	Early autumn legume sowing is needed to fix N. Legume cover crops can provide some N availability to following crops.	Buckwheat is better suited to spring sowing although autumn use in mixes is possible. Lupins typically need pH <7.	There is little published UK research in this area and further field information is needed.	

SOIL STRUCTURE

KEY TO SELECT A SUITABLE COVER CROP/MIX COMPONENT

Period for cover cropping: Autumn (to be followed by a spring crop)		Spring			Full season fallow		
Where is the main impediment?							
Shallow (c. 0-20cm)	Deep (c. 20-40cm)	Very deep (> 40cm)	Shallow (c. 0-20cm)	Deep (c. 20-40cm)	Very deep (> 40cm)	Shallow (c. 0-20cm)	Deep (c. 20-40cm)
Range of suitable types including cereals, brassicas, legumes and other broadleaf species.	Brassica cover crops (and possibly other deep rooted broadleaf cover crops).	Short duration cover crops are not well suited to this scenario.	Range of suitable types including cereals, brassicas, legumes and other broadleaf species.	Brassica cover crops (and possibly other deep rooted broadleaf cover crops).	Short duration cover crops are not well suited to this scenario.	Range of suitable types including cereals, brassicas, legumes and other broadleaf species.	Brassica cover crops and some legume species.
Suggested options/ingredients							
Oats, phacelia and brassicas (such as mustard or radish), but consider use of mixtures particularly if other depths are of interest.	Mustard or radish, but consider use of mixtures particularly if other depths are of interest.	None.	Cereals (rye or oats), legumes (e.g. trefoil, vetch and others) and other broadleaf crops (e.g. phacelia or brassicas).	Mustard or radish but consider use of mixtures particularly if other depths are of interest.	None.	Oats, legumes (such as trefoil or crimson clover), phacelia and brassicas (such as mustard or radish); potentially in mixtures.	Mustard, radish or possibly some clovers or other deep rooted species.
							Lucerne or Clover (perhaps sweet or red clover) may be suited (and potentially some brassicas).
Other comments (also see agronomy guidance pages for specific suggestions)							
Think about soil incorporation and rotational conflicts. Brassicas in particular can also reduce autumn nitrate leaching.	Consider brassica bolting and biomass issues. Brassica crops can also be very effective at reducing nitrate leaching.	Timeframe is too short.	Think about incorporation, rotational conflicts and crop duration.	Consider duration/rooting potential as well as seed set and volunteer risk.	Timeframe is too short.	Mixes including legumes would be relevant.	Research with legumes in this role is limited.
							While research suggests this approach has potential, there is little UK field data in this area.

SUMMARY TABLE

EFFECT OF COVER CROP TYPE

Cover crop	Crop type	Sowing (autumn)	Example sowing rates (single species)	Main uses/comment
Beans	Broadleaf (pulse)	Late Aug – Sept	100-200 kg/ha	Mainly used in fertility building as part of mixtures or single species. Better suited to later sowing than many legumes, but consider rotational conflicts.
Black medick (trefoil)	Broadleaf (legume)	August	8-10 kg/ha	Mainly used in fertility building mixes, faster growing than some clovers, and can improve soil structure. Consider rotational conflict with pulses.
Buckwheat	Broadleaf (polygonum)	August	70 kg/ha	Used around fertility building and particularly scavenging phosphorus. Buckwheat is not frost tolerant and is probably best used in mixtures.
Chichory	Broadleaf (Asteraceae)	August	15 kg/ha	Deep rooted cover crop (delivering soil structure benefits) better suited to longer term use especially where grazing is of interest. Can be used in mixtures.
Crimson clover	Broadleaf (legume)	August	10-15 kg/ha	Mainly used in fertility building mixes, faster growing than some other clovers, and can improve soil structure. Consider rotational conflict with pulses.
Lucerne	Broadleaf (legume)	August	20 kg/ha	Mainly used in fertility building mixes and can be better suited to droughty soils than some other legumes. Consider rotational conflict with pulses.
Mustard	Broadleaf (brassica)	Mid Aug – mid Sept	5-15 kg/ha	Competitive crop with benefits for soil around management of erosion, leaching and structure. Consider rotational conflict with oilseed rape.
Oats	Grass (cereal)	Mid Aug – mid Sept	30-100 kg/ha	Competitive crop with benefits around shallower soil management, leaching reduction and erosion mitigation. The sowing rate will depend on specific use.
Oilseed Rape (OSR)	Broadleaf (brassica)	Mid Aug – mid Sept	5-15 kg/ha	Competitive crop with benefits for soil around management of erosion, leaching and structure. Consider rotational conflict with oilseed rape.
Peas	Broadleaf (pulse)	Late Aug – mid Sept	200-400 kg/ha	Mainly used in fertility building as part of mixtures or single species. Better suited to later sowing than many legumes, but consider rotational conflicts.
Phacelia	Broadleaf (boraginaceae)	Mid Aug – mid Sept	c. 10 kg/ha	Competitive crop with benefits for soil around management of erosion, leaching and structure. Not entirely frost tolerant but unlikely to senesce fully over winter.
Radish	Broadleaf (brassica)	Mid Aug – early Sept	4-12 kg/ha	Competitive crop with benefits for soil around management of erosion, leaching and structure. Consider rotational conflict with oilseed rape.

Continued overleaf

Cover crop	Crop type	Sowing (autumn)	Example sowing rates (single species)	Main uses/comment
Ryegrass	Grass (Lolium)	Typically August - Sept	30-35 kg/ha	Competitive crop with benefits around shallower soil management, leaching reduction and erosion mitigation.
Sanfoin	Broadleaf (legume)	August	70 kg/ha	Mainly used in fertility building and grazing mixes but is less well suited to droughty soils than some other legumes. Consider rotational conflict with pulses.
Sweet clover	Broadleaf (legume)	August	10-15 kg/ha	Mainly used in fertility building mixes, quite slow growing but can improve soil structure (from longer residence). Consider rotational conflict with pulses and malting barley.
Vetch	Broadleaf (legume)	August - Sept	80 kg/ha	Quite a competitive legume and mainly fertility building mixes and can be later sown than some other legumes. Consider rotational conflict with pulses.
White clover	Broadleaf (legume)	August	10-15 kg/ha	Mainly used in fertility building mixes, quite slow growing but can improve soil structure (from longer residence). Consider rotational conflict with pulses.

The ingredients – cover crop components

Brassica types

Mustard

(*Brassica* and *Sinapsis* species e.g. *Sinapsis alba*).



Mustard is widely used and can be a competitive crop that grows rapidly, quickly covering the soil surface and developing a strong rooting system. The crop can also reduce nitrate leaching, help to manage soil erosion and improve soil structure.

Sowing: for autumn sowing mid August - early September is suggested (although conditions are more important than calendar date). Typically sown at c.10 kg/ha (range 5-20 kg/ha); but this varies with establishment conditions, method, mix used and seed size (thousand seed weights are often typically around 3-5 g). Seed costs vary substantially depending on mustard type but costs of £2-10/kg may be typical.

Management: Sowing depths and establishment are similar to oilseed rape, so mustard can be established either through broadcast or drilled routes depending on factors including seedbed conditions and equipment available. In a similar manner to oilseed rape, mustard cover crops should be managed for pigeons, slugs and flea beetles.

Typically, no other inputs are required through the growing period. Some growers favour using small amounts of starter nitrogen fertiliser, although recent Origins™, and other UK research, has only shown limited value to crop growth. Starter 'bagged N' should only be used where directed by FACTS qualified advice. Some mustard types will bolt more readily than others; as plant stems and other biomass develops it can become increasingly fibrous which can impact on destruction and breakdown. A range of mustard types are available and while impartial information on specific varieties/types is limited breeder/supplier information can be useful and help support selections.

Notes: Mustard is related closely to oilseed rape and other veg brassicas. This can impact on disease carry over, volunteers and other weed issues in the rotation. Where mustard is being used as part of a programme for improving soil condition, use in conjunction with soil amendments can be complementary. This section refers ostensibly to autumn sown mustard; in some instances mustard may also be a spring sown cover crop (although generally this is less common).

Radish

(various types, but usually *Raphanus* sp).



Radish types tend to be competitive and quick growing with strong deep tap roots. As cover crops, or components in mixtures, they can reduce nitrate leaching, help to manage soil erosion and improve soil structure (radish can be particularly deep rooted from an autumn sowing compared to other cover crops).

Sowing: for autumn sowing mid August - early September is suggested (although conditions are more important than calendar date). Typically sown at c.8 kg/ha (range 4-12 kg/ha); this varies with establishment conditions, method, mix used and seed size (thousand seed weights are often typically around 3-5 g). Seed costs vary substantially depending on type but costs of £4-10/kg may be representative.

Management: Sowing depths and establishment are similar to oilseed rape; so radish can be through broadcast or drilled routes depending on factors including seedbed conditions and equipment available. In a similar manner to oilseed rape, radish cover crops should be managed for pigeons, slugs and flea beetles (although some types are less susceptible than others). Often no other inputs are required through the growing period, although some growers favour using small amounts of starter nitrogen fertiliser. Recent Origins™ and other UK research has only shown limited value to crop growth and starter 'bagged N' should only be used where directed by FACTS qualified advice. Both rooting type (e.g. the size and nature of the tap root) and the propensity for bolting will vary with variety; but some radish types will stay as a vegetative rosette over the autumn, consequently radish can present less biomass to deal with in the spring compared to mustard. A range of radish types are available and, while impartial information on specific varieties/types is limited, breeder/supplier information can be useful and help support selections.

Notes: Radish is related closely to oilseed rape and other veg brassicas. This can impact on disease carry over, volunteers and other weed issues in the rotation. Where radish is used as part of a programme for improving soil condition, use in conjunction with soil amendments can be complementary.

Brassica 'sprinkles'

In addition to the key ingredients above there is a range of other 'sprinkles' (other brassica types) that could complement these ingredients; these may serve well as additions to mixtures or are perhaps suited for sole use in some circumstances. In general comments on sowing, management and rotational conflicts would be similar to those described previously.

Options are often selected around root characteristics e.g. stubble turnips (can be particularly useful where grazing provision is an objective) or tillage radish (which has a larger deep tap root) or trap/bio-fumigant activity (often used alone; see 'weed and pest management' section).

Legume species

Clovers

(various types, but often *Trifolium* sp).



Like all legumes, clovers have the ability to fix nitrogen (N) from the air and can also offer useful improvements to soil structure; collectively this can confer fertility building benefits. N is more likely to be fixed during spring rather than autumn growth, however from early sowing some nodulation and N fixing may be expected. From spring sowing levels of up to 150 kg/ha N pa are cited for a range of types. Clovers are commonly used in species mixtures.

Sowing: White clover, sweet clover and crimson clover can all be spring (March onward) or autumn (late July - mid August) sown. Where crops are autumn sown chances of establishment will diminish as sowing dates move toward September. Seed rates will vary but rates for white clover (*Trifolium repens*) (10 kg/ha, with thousand seed weights often around 0.5 g), sweet clover (*Melilotus officinalis*) (15 kg/ha, with thousand seed weights often around 2 g) and crimson clover (*Trifolium incarnatum*) (15 kg/ha, with thousand seed weights often around 2.5-3.5 g) could be considered representative for use as single

species. A range of other types of clover could also be sown. White clovers are often slow growing, but can be beneficial in mixtures and low input systems, crimson clover is potentially more suited to use in shorter (autumn sown) leys (or in mixes) and sweet clovers can produce a deep tap root if left for longer periods. Seed costs vary substantially with type, but costs of £6-10/kg may be representative.

Management: Clover species should generally be broadcast or drilled into firm, moist seedbeds at a shallow depth (a few mm) and rolled to improve establishment. White clover seed can be purchased as pelleted seed (with a clay coating to effectively increase the seed size). This can be useful and help to improve uniformity of distribution if seed is to be broadcast; some producers will recommend a slightly reduced seed rate here as well. As with most legumes, some pest protection can be required (e.g. beetles, weevils, slugs and pigeons) and inputs should be targeted as necessary. Depending on the type/mix being grown, topping (or grazing) may be beneficial during the season.

Notes: Clovers are legumes and as such rotational intensity needs to be considered where other legumes or pulses are grown in the rotation (e.g. for disease carry over). Clovers tend to be slow to establish and initially can be uncompetitive with weed species; however once established they will be more competitive. The N fixed by clover does not necessarily all become available to a following crop or at the time of requirement. Use of some mixtures of species can help with the fertility building benefits and some research suggests species such as crimson clover can be better suited to autumn ley use.

Black Medick

(*Medicago lupulina*, often referred to as yellow trefoil).



Black medick is a short-lived nitrogen (N) fixing annual legume with a low growth habit; it tends to be faster establishing than white clover (similar to crimson clover). This can offer both fertility building and soil structure benefits in autumn leys. Black medick has been used both alone and in mixtures and in research has performed relatively well from early August sowings.

Sowing: Black medick can be spring (March onward) or autumn (late July - mid August) sown; chances of good autumn growth decline as dates tend to September. Sowing rates vary with establishment conditions/method and seed size (with thousand seed weights often around 1.5-2.0 g), but c. 7-12 kg/ha are common where the species is used alone. Seed is cost likely to be c. £10-12/kg.

Management: Black medick is typically broadcast or drilled into firm, moist seedbeds at a shallow depth (a few mm) and rolled to improve establishment. As with most legumes, some pest protection and other inputs may be required (e.g. beetles, weevils, slugs and pigeons) but are not always necessary. In limited research at NIAB sites black medick has remained prostrate from autumn sowing; this has provided benefits around the following crop establishment and has not required topping over the autumn.

Notes: Black medick is a legume and as such rotational intensity needs to be considered where other legumes or pulses are grown in the rotation (notably for disease carry over). The N fixed by a legume does not necessarily all become available to a following crop or at a time of requirement; use of some mixtures of species can help in these situations. Black medick seems to be generally well suited for use as a component of mixtures.

Vetch

(*Vicia sativa*, also known as 'tares').



Vetch is a nitrogen (N) fixing annual legume that is commonly used in cover crop mixtures (although it could be used alone). There are a number of types of vetch available.

Sowing: Vetch can be spring (March onward) or autumn sown. For autumn sowing vetches can be sown later than clovers and medick species (August

through to early September). Seed is also larger than clovers (with thousand seed weights often of 40-80 g) and rates of 75-100 kg/ha are common when used alone. Cost is around £1-2/kg.

Management: Vetch needs to be deeper sown than clovers; possibly 2 cm into some tilth and moisture, consolidation can also be beneficial. Pest protection and other inputs may be required (e.g. for sitona weevils and sometimes pigeons), but is not always needed. Vetch can produce a lot of biomass for a legume cover crop but field accounts suggest it does not respond well to topping.

Notes: Rotational intensity should be considered where vetches are grown in rotation with other legumes. As with many legumes vetch is often used as a mix component and this can work well as the seed size and sowing requirements tend to be more in keeping with non-legumes compared to small seeded legumes. It should be noted, the N fixed does not necessarily become immediately available to following crops, but is typically retained in the soil system.

Legume 'sprinkles'

The legumes described previously are only a sub-set of those that could be used as key ingredients, and there are a range of possible 'sprinkles' that could be added to mixtures to complement these ingredients. Specific agronomic guidance on these additions will vary with crop type, although the rotational considerations would generally be common to all legumes.

Suggestion of other novel options to consider are:

Beans and peas

These can be a relatively cheap and farm available ingredients where there is frequently on farm production expertise. As larger seeded legumes they can be sown later than many other smaller seeded legumes, but appreciable N fixing is more likely to occur from earlier sowing. Autumn seeding rates of 100-400 kg/ha may be appropriate if used alone (lower end generally for beans and higher end for peas), but these options are perhaps most likely to be used within mixtures rather than as stand alone species.

Lucerne (Alfalfa)

This may be August sown as an autumn cover (often at c. 20 kg/ha as a single species). The crop will develop a tap root and can be suited to droughty soils. Over winter it often defoliates, but will generally recover. Lucerne usually needs to be destroyed in short-term leys and can require topping. Seed costs vary but are often around £10/kg.



Lucerne (Alfalfa)

Sainfoin

Traditionally a spring sown legume, but can be grown as a summer (August) sown legume (at c.70 kg/ha as a single species for autumn cover). The species tends not to like droughty soils, is highly frost tolerant (so will need to be destroyed in short term leys) and has excellent grazing potential (can be suited to use in longer leys and where it is not grazed it can need topping). Seed costs vary but can be around £2-3/kg.



Sainfoin field

Other options

Grasses: oats (*Avena* sp.), **rye** (*Secale* sp.), **rye-grass** (*Lolium* sp.) and others.



Graminaceous species typically give rapid ground cover from autumn sowings, often have relatively well understood agronomy and can be easy to grow compared to some other cover crop types (e.g. legumes). While growth is rapid and rooting can be vigorous, grass species are more likely to root more actively at shallow depths compared to those achieved with brassica roots (for example). However grasses are useful at opening up surface soil structures and commonly tend to be used in mixtures, but in some cases can be used alone.

Sowing: Oats and Rye: can be spring or autumn sown and seed rate will depend on conditions and use, but rates of c. 35 kg/ha (when used within autumn mixes for erosion management) up to 80-100 kg/ha (where stands are needed as a single species for example for weed competition) are common; seed costs are often around £1-2/kg and thousand seed weights of 30-50 g can be representative.

Italian (and Perennial) ryegrass can be sown in April or August/September with seed rates (from thousand seed weights often around 1.5-2.5 g) typically around 30-35 kg/ha (perennial ryegrass is suited to longer term stands); seed costs are £2-4/kg. Westerwolds rye-grass can be drilled later than other rye-grass, potentially into October for cover crop use, and is again typically drilled at 25-35 kg/ha in the autumn; seed costs are £2-4/kg.

Management: Oats and rye can be established via drilling (at a similar depth to cereal crops) or through broadcasting and rolling routes, provided there is adequate tilth. Ryegrass should generally be drilled into firm seedbeds at 1-2 cm and rolled to improve establishment. Post establishment, frequently no other inputs are required through the growing period, however, sites may need to be managed for pest damage should this occur. Ryegrasses can provide grazing opportunities or may need to be topped during the autumn.

Notes: A range of oat types in particular are available and while impartial information on specific types is limited, breeder/supplier information can be useful and help support selections. Practically many farmers using oats will often use a spring oat variety in the autumn. While consideration should also be given to carry over or increased disease pressure to cereal crops, volunteer management in following crops is likely to be the greater immediate practical conflict. In this respect the management of volunteer oats, in particular, can often be less problematic in following spring crops than many other cover crop choices.

Phacelia

(*Phacelia tanacetifolia*)



Phacelia is fast to establish, highly competitive and the purple flowers provide a source of nectar and seeds beneficial to insects and other wildlife.

Sowing: For spring sowing Phacelia can typically be sown March onwards, but it is also suitable for early autumn sowing (e.g. a similar window to oilseed rape of mid August to early September). Seeding rate varies with establishment conditions/method and seed size (with thousand seed weights of 2-3 g), but c.8-10 kg/ha is suitable where it is used alone; seed costs vary but £8-10/kg is typical.

Management: Phacelia can be shallow drilled (perhaps 1-2 cm) or broadcast to suit conditions and equipment. Frequently no other inputs are required through the growing period, however, sites may need to be managed for pest damage should this occur. Autumn sowings do not tend to get overly tall and topping is not usually required. Some canopy management/topping could potentially be required from spring sowing to prevent seed set in certain scenarios. From autumn sowing Phacelia is not fully frost tolerant, but the degree of winter senescence depends very much on the season and crop destruction is often needed.

Notes: Phacelia is a boraginaceae and unrelated to most other crops grown commonly in UK rotations (although is related to borage, echium, buglossoides etc). Unless seed set is an objective (e.g. for environmental goals) growers should seek to remove the crop before seeds set to limit volunteer problems in following crops. Should Phacelia set seed there can be a requirement to manage volunteers in following crops, although this is more likely to be an issue where herbicide choice is limited. NIAB TAG research has indicated positive, but variable, yield responses in spring barley crops following phacelia.

Other 'sprinkles'

There are a range of options that can be used as cover crops and finding species that are outside legumes, brassicas or cereals can be a good way to minimise rotational conflicts. In addition to the key ingredients described previously there are several possible 'sprinkles' that could be added to complement seed mixtures. Two possible options to consider are as follows:

Buckwheat

(a Polygonaceae)

This species typically does better from spring establishment, but can be summer sown (August); buckwheat can be useful at scavenging phosphorus (potentially improving availability), but is short-lived and not frost tolerant. Buckwheat is probably best used as a component of mixtures. Seed rates are up to 70 kg/ha (used as a sole species) and typical costs are often £2-4/kg.

Chichory

(an Asteraceae)

Can be spring (March onwards) or autumn sown (typically August). The species produces a very deep and vigorous tap root, but is slow growing and is often suited to longer term leys, but can be used as a component of mixtures. Chicory is frost tolerant and can fit well with grazing use. Seed rates are up to 15 kg/ha (used as a sole species) and typical costs are often £10-15/kg.



Buckwheat (a Polygonaceae)

The mixing bowl – combinations of components in a cover crop

There are no hard and fast rules about selecting cover crop mixtures and the selections should be governed by the primary objective of the cover crop as well as other factors including budget, potential rotational considerations and farm equipment/scenario. In some cases this may require a bespoke mix or the generation of simple mixes on farm through the collection of various ingredients, but in others, pre-formulated mixes sold through a range of seed merchants will serve well.

For further information formulating and growing mixes see the section 'Cooking; growing cover crops' and consult seed supplier or other specialist guidelines.

The suggested mixes presented in this section are some that may be used commonly on farm, have

worked well in field research or have been suggested through consultation. However, these are only examples and growers are encouraged to experiment with mixes suited to their own goals and situations; why not try some field strips of different cover crops or some split field comparisons.

The following are simple mixes ostensibly for use in soil management (e.g. erosion, leaching reduction or soil structure situations). However, these mixes would also potentially provide other benefits around green manure use, habitat provision and weed competition. These mixes could also be extended to include other species to provide a wider range of root types (e.g. try oil and tillage radish with oats rather than just a two way mix) or add other species to widen objectives (e.g. consider adding in vetch as a fertility building legume).



Cover crop mixes

Oilseed rape and oats

(use: soil management, particularly erosion mitigation)

Tends to be used as a relatively inexpensive mix to manage soil erosion (providing vigorous, effective ground cover at key periods); although will also provide some soil conditioning, nutrient retention and green manure benefits.

Seed rates: will vary with circumstance but rates of around 30-50 kg/ha for the oats and 3-10 kg/ha for the oilseed rape can be suited to a range of situations.

Sowing dates: typically mid August to mid September (depending on conditions).

Complementary species: this simple two way mix could be extended by adding in other components. For example, addition of vetch or crimson clover would introduce a faster growing legume for autumn sowing. In addition rye could be used instead of oats.

Other comments: use of oilseed rape as a cover crop and closer brassica rotations are a common cause of concern with this mix.

Radish and oats

(use: mainly for improving soil structure and erosion management)

The root depths and structures of the two species complement each other well and can be useful at developing soil bio-pores over a range of depths. This simple competitive mix will also provide useful erosion mitigation, nutrient retention and green manure benefits.

Seed rates: will vary with circumstance but rates of around 30-50 kg/ha for the oats and 3-8 kg/ha for the radish can be suited to a range of situations.

Sowing dates: typically from mid August to early September.

Complementary species: this two way mix could be extended by adding in other components. For example, phacelia would add a third species with a different rooting habit (to aid soil structuring) or legumes such as vetch or crimson clover would introduce a nitrogen fixing species. In addition, rye could be used instead of oats.

Other comments: closer brassica rotations are a common cause of concern with this mix.

Phacelia and oats

(use: potentially suited to soil conditioning and improving structure)

Where additional brassica use in a rotation is a concern phacelia can also be a vigorous rooter and the combination with oats again would give complementary root depths.

Seed rates: there is less field experience with this mix, but rates of 30-50 kg/ha for the oats and 3-6 kg/ha for the phacelia may be suited depending on circumstance and goal.

Sowing dates: typically from mid August to early September.

Complementary species: as with previous mixes other components such as this two way mix could be extended by adding in other components. For example, legumes such as vetch or crimson clover to added (to introduce nitrogen fixing species). In addition, rye could be used instead of oats.

Other comments: the mix will not conflict with brassica crops in the rotation, but equally probably will not root to the same depth as brassica species.

The following are simple mixes ostensibly for fertility building and include legumes to fix/capture nitrogen in a way that is potentially more available to the crop. These mixes could also provide wider benefits around habitat provision, leaching reduction and soil condition improvements. In addition to the suggested mixes also consider simple approaches, such as the use of beans as a cover crop (or part of the mix) or more adventurous approaches with multi-way mixes (e.g. potentially adding other ingredients to improve deep soil conditioning).

Black medick and Phacelia

(use: mainly for habitat (insects) and fertility building benefits)

This mix is a combination of an active rooting phacelia cover and one of the faster developing, nitrogen fixing, legume species. Phacelia is also very attractive to bees and other pollinators.

Seed rates: as a guide seed rates of around 3-5 kg/ha for the medick and 3-8 kg/ha for the phacelia, depending on circumstance, may be suited to this mix.

Sowing dates: typically from mid to late August (the legumes tends to do less well from September sowing).

Complementary species: other choices of N fixing legume species for use in the mix, either as well as or instead of black medick, could include vetch, crimson clover or beans.

Other comments: this mix provides an option that will not conflict with brassica crops in the rotation, although medick is a legume so the proximity of pulse crops in the rotation should be considered. These components have performed well individually in research trials, but there is a need for more field experience and data with the mix.

Black medick (or other legume) and oats

(use: mainly fertility building benefits)

This mix is combination of the vigorous root and canopy development of oats and options for the faster developing, nitrogen fixing, legume species.

Seed rates: as a guide seed rates of around 3-5 kg/ha for the medick and 30-50 kg/ha for the oats. This mix will depend on circumstance and goal.

Sowing dates: generally from mid to late August (the legumes tends to do less well from September sowing).

Complementary species: other choices of N fixing legume species for use in the mix, either as well as or instead of black medick, could include vetch, crimson clover or beans. In this mix crimson clover (possibly at 5-8 kg/ha) would perform similarly to black medick and is also a relatively faster growing legume (albeit

with a different growth habit), but field observation suggests it may be more attractive to certain pollinators. The use of vetch, perhaps at 20-40 kg/ha, is another legume option which may cope better with slightly later sowing. Beans or peas could also be added as a relatively cost effective legume mix component. Alternatively a multi-way mix with several of these components could also be considered.

Other comments: this mix provides an option that will not conflict with brassica crops in the rotation, although medick is a legume so the proximity of pulse crops in the rotation should be considered. The black medick and crimson clover components have performed well individually in trials and there is some expert opinion to support the mix, but there is a need for more field experience and data.

The following are a couple of examples of possible wider mixes. These suggestions are by no means exhaustive but are illustrative of potential multiway mixes.

Legume species mixture

(use: mainly fertility building but also soil condition benefits)

The mix (white clover, black medick, lucerne and crimson clover) is based on one developed and tested in Defra LINK project LK09106 (using legume-based mixtures to enhance the nitrogen use efficiency and economic viability of cropping systems). This mix has been used by NIAB TAG and others and estimates of return from research vary, but margin over nitrogen improvements of around £65/ha have been recorded in NIAB trials.

Seed rates: the total rate tends to be 10-15 kg/ha with rate ranges for the components as follows: white

clover (2-4 kg/ha), black medick (trefoil) (2-4 kg/ha), lucerne (3-5 kg/ha) and crimson clover (3-5 kg/ha).

Sowing dates: generally in early to mid August (and tends to do less well from a September sowing).

Other comments: while this legume mix provides an option that will not conflict with brassica crops in the rotation the proximity of pulse crops in the rotation should be considered. Due to the range of seed sizes in the mix, care should be taken to ensure uniformity of seed distribution during sowing.

Radish, crimson clover, oats and buckwheat

(use: mainly for improving soil structure and nutrient availability)

The root depths and structures of the species complement each other well and can be useful at developing soil bio-pores over a range of depths. In addition there is some suggestion from research that buckwheat can be effective in sequestering phosphate. This competitive mix will also provide useful erosion mitigation.

Seed rates: will vary with circumstance but rates of around 30-50 kg/ha for the oats and 3-8 kg/ha for the radish and crimson clover and (perhaps) 10-20 kg/ha of buckwheat.

Sowing dates: typically from early August to early September.

Complementary species: if a legume was not required in the mix phacelia could be added as well as or in place of the crimson clover. In addition rye could be used instead of oats.

Other comments: close brassica and pulse rotations could be a cause of concern with this mix.



Phacelia mustard mix

‘Cooking’ – growing the cover crop

Cover crop selection

There are few hard rules with selecting cover crops, but guidance on specific choices either to use alone or in mixture begins on page 14.

Seed mixtures

The use of mixtures is common. There are a wide number of pre-generated commercial cover crop seed mixes available, but equally growers could develop their own. While there is a need for further comparative data on mixes, the suitability of mix components and field experience can be a useful guide to their value.

The mixes and approaches suggested in this guide are provisional and based on farm practice, limited experimental data and discussion with growers. When selecting cover crops to use in mixtures, consider how the components help to achieve the overall objectives and complement each other; for example is one component likely to out-compete another, are the lifecycles, breakdown characteristics etc suitably matched or do the functional traits help to achieve the overall goal of the mix.

There are also some practical aspects to consider around using mixtures. For example, when seed sizes

differ appreciably in mixes there can be a need to mix the seed periodically in the drill to avoid separation. In addition some concession on drilling depth can be needed when mix components have markedly different requirements. In general it is advisable to select components that minimise these differences, although this is not always possible and other solutions (such as separately broadcasting and drilling some mix elements) can be used if needed.

Sowing date

This will be guided by the specific components of a mix, but broadly autumn sown mixes tend to be sown in early August - mid September. Most legumes need to be earlier in this window (by mid August) but other cover crops (e.g. brassicas and cereals) can often be sown later. In general, earlier sowing tends to show greater growth over the autumn, however, good soil/seedbed conditions are more important than specific date. For many cover crop species, seed should ideally be established into surface tilth in a firm, moist seedbed and rolled to improve moisture conservation and establishment. The relationship between sowing date and autumn cover crop growth from Origins™ farmers in autumn 2014 is presented in Figure 2 while some examples of autumn growth (measured as Green Area Index (GAI)) are presented in Figure 3.



Figure 2. The impact of sowing date on mean autumn cover crop green area index (GAI) in Kellogg's Origins™ sites from assessments made in October 2014.





Figure 3. Autumn (October) and spring (February) cover crop green area index (GAI) in selected sites over the Kellogg's Origins™ programme (2014). Sowing dates were site 1 (07/08/14), site 2 (31/08/14) and site 3 (23/08/14). Green Area Index (GAI) is a measure of the ratio between the total area of all green tissues and the area of ground from which they come.



Sowing rates

Specific seed rates will depend on the inclusions and specific mix. Further information on seed rates can be found on page 12, but as a broad steer in multi-way mixes, it is often common to sow at less than full rate. As a guide perhaps quarter to up to a half of the

rate (cf. that would be used for that cover crop grown alone) would be typical of the main components in several mixes; but this will also be guided by the objective, the number of components in the mix and the overall budget for the cost of the mix.

Establishment

The method of establishment varies with cover crop type, equipment available, field conditions, soil type and farm scenario; however, generally, a cover crop is drilled or broadcast followed by seedbed consolidation. Establishment with broadcasting (possibly trash raking to distribute straw and generate tilth, followed by broadcasting and rolling) tends to be cheaper, but can be more variable and seed distribution uniformity should be considered. When drilling seed mixes, a range of drill types can be used; single pass systems are often adopted to improve timeliness and reduce cost. Thought should also be given to row width/seed spread; the 'floating' drill method (either over or just engaging with soil) has been employed by a number of growers.

Management and inputs

This will be guided by the specific mix components. While monitoring autumn sown cover crops for input requirements is recommended, in some situations there may be no need for any inputs. However, in some situations pest protection could be needed for grazing by pigeons, slugs or insect pest damage.

The need for autumn starter nitrogen (N) fertiliser is sometimes advocated. This was assessed on two Origins™ sites in 2014; use resulted in some growth increase, but differences were generally small (mean of 10-20% increase, Table 5). While there was some interaction with cover crop type, the growth benefits appear to be limited from this data. Starter N also tended to increase autumn weed populations. By the spring differences due to starter N were difficult to detect.

Table 5. Mean data from 15 cover crop comparisons with, and without, starter fertiliser carried out over two sites in the Kellogg's Origins™ programme during 2014/15.

Comparison of plant counts (per m²) and green area index (GAI; as a measure of canopy size).

Treatment	Autumn assessment			Spring assessment		
	Cover crop		Weed	Cover crop		Weed
	Count (plants/m ²)	GAI	Count (plants/m ²)	Count (plants/m ²)	GAI	Count (plants/m ²)
Without N	61	1.6	24	29	1.1	21
With N	60	1.8	51	30	1.2	27

Cover crop destruction/following crop establishment

Methods of destruction vary markedly and will depend on soil type, growth (canopy size and type of growth), available equipment and objectives. Autumn-established covers can be killed off by frost action, grazing/destroyed mechanically or sprayed off with glyphosate early in the year (while other herbicides could be used glyphosate tends to be used most commonly).

Where cover crops are being used to help outcompete/manage pernicious weeds, more than one glyphosate application may be needed (as the cover may shelter some weeds). This is often followed by some degree of spring incorporation (ideally via non-inversion systems) prior to establishment of the following crop, although in other situations, single pass drilling through residues is used.

The interaction of soil type and sowing system (e.g. drill type) will also have a bearing on destruction/method and in general options and opportunities are more extensive on light to medium soils, whereas on heavier soils thought should be given to opening up cover crops early enough to allow the soil surface to dry (particularly where surface structure/drainage is poor, surface drying can be needed on heavier soils ahead of drilling). The specific approach used is likely to be highly farm specific; when selecting a cover crop mix it can be useful to visualise what it might look like in the spring and consider how this will integrate with your proposed destruction method and drilling equipment. Where bio-fumigant or trap crops are sown there are usually specific destruction requirements in order to attain the full benefits.

Rotational planning

Potential rotational conflicts should be considered with any cover crops or cover crop mixture. For example the likely risks and impacts of shortening brassica and pulse rotations or likely issues with volunteer management in following crops.

Field research on the longer term impacts is limited, but for brassicas NIAB TAG's New Farming Systems research, over an eight year period, shows some reduction (c. 6%) in oilseed rape yield associated with short (alternate) rotations of brassica cover crops, although this is less than would be expected from a short (alternate) oilseed rape rotations (c. 12%). The research also suggests there may be some potential to further mitigate these losses if shallow non-inversion systems were used (possibly better management of cover crop volunteers). Problems with specific pathogens such as clubroot and verticillium were not apparent in this study, but could be of concern on some sites (e.g. those with a history, where local pressure is high or where field conditions are suited).

Information on the impact of legume inclusions in pulse crops is very limited and will be influenced by the pathogens in question, the legumes grown, the farm site and environmental conditions, but current thinking suggests at least a three year break before a grain legume crop may be appropriate.

Use of cover crops in EFAs

Cover choice is generally best guided by the end objective and the farming scenario, but those wishing to grow catch crops and cover crops as an Ecological Focus Area (EFA) option will need to follow a more predefined set of options. Further details can be found in the relevant Defra CAP guidance; essentially the guidance sets out a predefined list of options, suitable mixtures and key dates that need to be followed. While cover crop use within this mechanism will contribute to CAP greening, those wishing to grow cover crops outside this mechanism will have greater flexibility.

'The new Common Agricultural Policy schemes in England: October 2014 update' (page 11) states:

Growing catch crops and cover crops as an EFA

On page 30 of our August update, we said we would publish more information about which catch and cover crops will count as an EFA. Catch and cover crops are designed to protect the soil and use available nutrients between harvest and sowing. The crops farmers can grow as an EFA are those that give the best chance of:

- establishing within the sowing period
- growing quickly
- achieving ground cover
- having different rooting depth types.

Farmers must use a sown mix of at least two different cover types (one cereal and one non-cereal). However, grass can be used as either a catch crop or a cover crop as long as it was undersown in the previous crop and is sufficiently established. Crops that farmers can grow in the sown mix are:

- Rye
- Vetch
- Phacelia
- Barley
- Mustard
- Oats
- Lucerne

This list is based on the crops that have been used successfully for Environmental Stewardship in recent years. The regulations don't allow farmers to include crops that are usually grazed, so we haven't included kale or stubble turnips. Using crops from this list will give the soil surface the best chance of protection from erosion. It will also help to make sure that available nutrients are taken up by the plants. Once the catch/cover crop is destroyed, farmers should take care to ensure that all those benefits are not lost. So, they should avoid grazing and establish the next crop quickly. Farmers can include other crops in their catch crops or cover crops, but these areas cannot count as an EFA.

Further information can also be found in the CAP updates at www.gov.uk and search for Common Agricultural Policy (CAP) Reform or scan the QR code.



Seed sourcing

There are a wide range of companies providing cover crop seed, with no particular preference these would include (among a wide range of others):

Boston Seeds: www.bostonseeds.com

Green Manure: www.greenmanure.co.uk

Bright Seeds: www.brightseeds.co.uk

Kings: www.kingscrops.co.uk

Cotswold Seeds: www.cotswoldseeds.com

Pearce Seeds: www.pearceseeds.co.uk

DSV: www.dsv-uk.co.uk/cover-crops

Further information

Cotswold Seeds; 'Sort out your soils' available through the following link:
www.cotswoldseeds.com/files/cotswoldseeds/Cotswold_Green_Manures_final.pdf

Döring TF, Baddeley JA, Brown R, Collins R, Crowley O, Cuttle S, Howlett SA, Jones HE, McCalman H, Measures M, Pearce DB, Pearce H, Roderick S, Stobart R, Storkey J, Tilston EL, Topp K, Watson C, Winkler LR, and Wolfe MS (2013). Using legume-based mixtures to enhance the nitrogen use efficiency and economic viability of cropping systems. HGCA Project Report 513.

'Managing Cover Crops Profitably' from the U.S. Department of Agriculture.
See: www.mccc.msu.edu/documents/ManagingCCProfitably.pdf

Stobart RM and Morris NL, (2013), Approaches to cover cropping and the impact on soils and farming systems, Aspects of Applied Biology 121 (Rethinking Agricultural Systems in the UK), pp43-50.

Stobart *et al* (2015), Developing the use of cover crops on farm through the Kellogg's Origins™ grower programme, Aspects of Applied Biology 129 (Getting the Most out of Cover Crop), pp 27-34

Morris NL, Stobart RM and Orson JH, (2014), An appraisal of research, best practice and communication approaches for the management of soil structure, Felix Cobbold Trust review (see the member area of the NIAB website).

Through the ARTIS cover crop e-learning training programme: www.artistraining.com/e-learning

It is intended that the information provided in this document will continue to evolve and your feedback and suggestions for improvement are welcomed. Please email nac@niab.com or origins@kellogg.com with comments for the further development of this publication.

Acknowledgements

Thanks are extended to:

Kellogg's, farmers and other specialists participating in the Kellogg's Origins™ programme.

The Morley Agricultural Foundation (TMAF) and the JC Mann Trust for their support of the New Farming Systems cover crop work.

The NIAB National Agronomy Centre Initiative research; supported by TMAF.

Partners in Defra LINK project LK09106 (Using legume-based mixtures to enhance the nitrogen use efficiency and economic viability of cropping systems).

Other commercial sources of information including Cotswold Seeds, Kings, RAGT and other cover crop researchers.



NIAB TAG, Huntingdon Road, Cambridge CB3 0LE, UK
Tel: +44 (0)1223 342200 info@niab.com  @niabtag
niab.com

While every care has been taken in the preparation of the advice contained in this booklet, NIAB TAG and Kellogg's cannot accept responsibility for any loss of inconvenience arising from following the information herein.

© 2015. All information produced by NIAB TAG is protected under copyright and is not to be reproduced in any form or distributed to other parties without prior written permission of NIAB TAG.

Design and produced by: Cambridge Marketing Limited, www.cambridgemarketing.co.uk